

2021

BREAST SEMINAR SERIES

Faculty

LÁSZLÓ TABÁR, MD, FACR (Hon) Course Director  
*Professor emeritus of Radiology*

Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach

AN INTERACTIVE,  
UNIQUE LEARNING EXPERIENCE  
LIVE WEBINAR

*Aug 21 and Aug 28*

*and*

*Sept 04 and Sept 11*


NEW  
course  
design

*Designed for:*

**Radiologists • Surgeons • Pathologists  
Gynecologists**

*This course provides extensive knowledge about diagnostic breast imaging, differential diagnosis of breast diseases, implications for management and newest diagnostic technologies*

3D image of the breast tissue



3D image of the breast tissue

**8 Category I  
CME credit hours.**



2021  
BREAST SEMINAR SERIES of MEI

Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

**László Tabár, MD, FACR (Hon)**  
*Course Director*

## FACULTY



**László Tabár, MD, FACR (Hon).**  
Course Director

*Professor emeritus of Radiology, Uppsala University, Sweden*



Photographs from the collection of the non-profit Tabar Foundation dedicated to Research and Education for Breast Cancer




2021

BREAST SEMINAR SERIES of MEI

Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

**László Tabár, MD, FACR (Hon)**  
*Course Director*



Mammography Education, Inc. is accredited by the Accreditation Council for Continuing Medical Education to sponsor continuing medical education for physicians. Mammography Education, Inc. designed these medical education activities for a maximum of **8 credit hours in Category I** of the Physicians' Recognition Award of the American Medical Association. Each physician should claim only those hours of credit that he / she actually spent in the educational activity.

## NEW COURSE DESIGN

- \* The lectures on each major subject will be followed by **immediate feedback** and discussion.
- \* During the course the attendees will progressively **improve their interpretive expertise**, as they learn all important findings explained with the help of large format thin section and 3-dimensional histology images.
- \* These skills will lead to greater confidence in analyzing and interpreting microcalcifications on the mammogram.
- \* Special emphasis will be placed on **finding early phase breast cancers**.
- \* All abnormal cases are fully worked up and the **complete imaging workup will be presented in detail, including ultrasound, MRI and large section histopathology**.






8:00 AM PDT (California daylight saving time). 10 AM CDT (Chicago). 11:00 AM EDT (New York). 5:00 PM CET (Central European Time, Stockholm)

CONTENT OF THE INTERACTIVE SESSIONS - SPREAD OUT IN FOUR DAYS


- Evaluation of asymmetric densities, specific for normal fibroglandular tissue. Detection and workup of non-specific asymmetric densities without architectural distortion. Parts I and II.
- Detection and didactic workup of benign and malignant non-specific asymmetric densities with architectural distortion. Parts I and II.



Radial scar






Neoductogenesis (DAB)



Diffusely infiltrating cancer of mesenchymal origin

ANALYSIS of **BENIGN RADIATING STRUCTURES** on the mammogram, originating in the ducts:  
**Radial scar / sclerosing ductal hyperplasia**







2021  
BREAST SEMINAR SERIES of MEI


László Tabár, MD, FACR (Hon)  
Course Director

Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

ANALYSIS of **MALIGNANT LESIONS** PRESENTED as non-calcified RADIATING STRUCTURES on the mammogram. Clinical presentation, mammographic appearance and outcome.

- **Duct forming invasive carcinoma / Neoductgenesis** cases presenting on the mammogram as architectural distortion. **The role of MRI in diagnosing diffuse breast cancer.**

**Interactive session for detecting architectural distortion on the mammogram.**



Non-calcified architectural distortion: extensive duct forming invasive cancer

László Tabár, MD  
Tibor Tót, MD, Peter B. Dean, MD

**In 3D**

Breast cancer originating from the major ducts

**Ductal Adenocarcinoma of the Breast (DAB), Part 7**

Architectural distortion on the mammogram without calcifications or nipple discharge

Mammographic-MRI-subgross (3D) histologic correlation of this extensive micropapillary cancer originating from the major ducts presenting as architectural distortion.

Architectural distortion on the mammogram without calcifications or nipple discharge

**In 3D**

Printed in China  
ISBN: 978-0-988541-9-8  
9 780988 541908


There are two main groups of diffuse breast cancers presenting on the mammogram as large regions of architectural distortion; these account for about 25% of all breast cancers and tend to have a poor outcome: 1) **Neoductgenesis**, i.e. "duct forming invasive carcinoma", the topic of this volume, often erroneously diagnosed as "DCIS", and 2) **Diffusely infiltrating breast cancer**, the topic of Vol. XI.

This volume demonstrates the DAB subgroup where the unimaturally high concentration of abnormal, tumor-filled ducts results in an asymmetric density with architectural distortion on the mammogram and often causes a palpable "thickening". Detecting architectural distortion on the mammogram and diagnosing the underlying disease correctly is a challenge for the radiologist. Breast cancers originating from the major ducts (DAB) are characterized by the formation of new, duct-like structures through the process of Neoductgenesis.





ANALYSIS of MALIGNANT LESIONS PRESENTING as RADIATING STRUCTURES on the mammogram. Clinical presentation, mammographic appearance and outcome, cont.


2) Diffusely infiltrating breast cancer of mesenchymal origin: the most deceptive and frequently missed cancer of the breast. The value of ultrasound and MRI in finding and diagnosing this spider's web-like malignancy. Case demonstrations, large section histopathologic-imaging correlation. Long-term outcome.



Example 1. Multimodality workup of a huge diffusely infiltrating breast malignancy of mesenchymal origin.



Example 2. Diffusely infiltrating (spider's web-like) carcinoma of mesenchymal origin in the upper half of the breast and a spherical, round lesion, originating from the TDLU (AAB) is seen in the lower portion of the left breast.



Interactive session for detecting architectural distortion on the mammogram.



2021  
BREAST SEMINAR SERIES of MEI

**László Tabár, MD, FACR (Hon)**  
*Course Director*

Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

**For more information and  
registration please contact:**

**Mammography Education, Inc.  
4429 E. Spur Drive  
CAVE CREEK, AZ 85331, USA**

Phone: (480) 419 0227

Fax: (480) 419 0219

e-mail: [info@mammographyed.com](mailto:info@mammographyed.com)

Internet: [lectures.mammographyed.com](http://lectures.mammographyed.com)

THE SCHEDULE IS SUBJECT TO CHANGE WITHOUT NOTICE AND DOES NOT REPRESENT A COMMITMENT ON THE PART OF M.E.I.  
ALL RIGHTS RESERVED INCLUDING THE RIGHT OF REPRODUCTION IN WHOLE OR IN PART OF ANY FORM.

VISIT US ON THE INTERNET: [HTTP://WWW.MAMMOGRAPHYED.COM](http://WWW.MAMMOGRAPHYED.COM) COPYRIGHT ©



A photograph from the collection of the non-profit Tabar Foundation dedicated to Research  
and Education for Breast Cancer





2021

BREAST SEMINAR SERIES of MEI



Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

**László Tabár, MD, FACR (Hon)**  
*Course Director*

## Breast Cancer Early Detection with Mammography

Casting Type Calcifications: Sign of  
a Subtype with Deceptive Features


László Tabár  
Tibor Tot  
Peter B. Dean



## Breast Cancer Early Detection with Mammography

Crushed Stone-like Calcifications:  
The Most Frequent Malignant Type

László Tabár  
Tibor Tot  
Peter B. Dean




[www.thieme.com](http://www.thieme.com)

## Breast Cancer The Art and Science of Early Detection with Mammography

László Tabár  
Tibor Tot  
Peter B. Dean



Calcification,  
Microcalcification,  
Histopathologic Correlation




## Teaching Atlas of Mammography

László Tabár  
Peter B. Dean

With the contribution of Tibor Tot

4th edition






2021  
BREAST SEMINAR SERIES of MEI


Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

László Tabár, MD, FACR (Hon)  
Course Director


László Tabár, MD  
Tibor Tot, MD, Peter B. Dean, MD



## Understanding the Breast in Health and Disease



In 3D



Multifocal breast cancer



Sea urchins

In 3D



In situ ductal carcinoma




Banana flower


The basic structural elements of the female breasts are illustrated here in true 3-dimensional (3D) images and described in this Volume I by three breast cancer experts with decades of experience in the diagnosis of breast diseases. These images provide the best way to understand the great variability of the normal breast structure and the changes brought about by benign and malignant diseases.

[www.mammographyed.com](http://www.mammographyed.com)

László Tabár, MD,  
Tibor Tot, MD, Peter B. Dean, MD,  
Miklós Tarján, MD




cysts in a prostate




breast cysts


## Prostate and Breast: Brother and Sister Organs



In 3D




Prostate calcifications



Laminated calcifications  
in the prostate



In 3D



Laminated calcifications in  
the breast



Rowan berries

Even as the risk of getting prostate and breast cancer is rising, early detection through screening and treatment in an early stage are significantly lowering the risk of dying from these diseases. This series of 3D books aims to empower both men and women with knowledge about their health. Although all of us are at risk of developing cancer or less serious problems in one or the other of these two organs, education will help us seek the benefits provided by modern health care and expect excellence from health care providers.






2021  
BREAST SEMINAR SERIES of MEI

László Tabár, MD, FACR (Hon)  
Course Director


Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

*László Tabár, MD*  
*Tibor Tot, MD, Peter B. Dean, MD*



Breast cancer of ductal origin with microcalcifications

## Ductal Adenocarcinoma of the Breast (DAB), Part 1



**In 3D**



8 mm poorly differentiated invasive breast cancer associated with neoductgenesis (DAB)

A photograph reminiscent of neoductgenesis with associated tiny invasive tumors



**In 3D**




Fragmented casting type calcifications make the cancerous duct-like structures visible on the mammogram.

Neoductgenesis is a frequent phenomenon in the plant world


The mammogram is a true representation of the structural changes induced by the genetic constellation of each breast cancer subtype. The mammographic/MRI/ultrasound presentation of a particular subtype reflects the nature and extent of the underlying disease process, and when correctly interpreted, can guide patient management and help in predicting the long-term outcome. This information is available at the moment of diagnosis, without the additional expense and time necessary for molecular and immunohistochemical analysis.

*László Tabár, MD*  
*Tibor Tot, MD, Peter B. Dean, MD*




An axillary lymph node populated with metastases mimicking *in situ* cancer

## Ductal Adenocarcinoma of the Breast (DAB), Part 2




**In 3D**



Metastases within an axillary lymph node mimicking cancer *in situ*

**In 3D**




Stereoscopic image pair of the DAB with calcifications within a lymph node

Breast cancers originating from the major milk ducts (breast cancer of ductal origin, DAB) occasionally cause axillary lymph node metastases which are similar in appearance at histology to DAB in the breast. Regardless of whether or not the myoepithelial cell layer is demonstrable, the decisive question is how do the duct-like structures grow inside the lymph nodes? Although the histopathologic appearance will be termed by pathologists as invasive cancer, i.e., when found in the prostate or in the axillary lymph node(s), a similar histopathologic appearance is termed "DCIS" when found in the breast. In reality, we face "duct forming invasive cancer" with poor outcome (neoductgenesis) in the breast, in the prostate and in the axillary nodes.






*László Tabár, MD*  
*Tibor Tot, MD, Peter B. Dean, MD*




Micropapillary breast cancer of ductal origin associated with a normal TDLU

### Ductal Adenocarcinoma of the Breast (DAB), Part 3




In 3D



Neoductgenesis (DAB) associated with angiogenesis


In 3D



Normal atrophic ducts and cancerous, distended ducts side by side


Breast cancers that originate in the major milk ducts (ductal adenocarcinoma of the breast, DAB) are diffuse and often extensive. The disease may occupy an entire lobe from the nipple to the chest wall, and frequently extends close to the skin. For these reasons, breast conserving surgery and skin or nipple sparing mastectomy of DAB cases carry a higher risk of local/regional/distant recurrence. In addition: 1) a considerable portion of the disease may lack calcifications, often occur for the imaging methods. 2) This subtype of breast cancer is less responsive to postoperative radiotherapy.

*László Tabár, MD*  
*Tibor Tot, MD, Peter B. Dean, MD*




Paget's disease of the nipple      Paget's cells in the epidermis of the nipple

### Ductal Adenocarcinoma of the Breast (DAB), Part 4




In 3D



Paget's disease of the nipple and breast cancer of ductal origin

In 3D



Cancer-filled duct in Paget's disease with angiogenesis

One of the features which is unique to breast cancers originating from the major ducts (DAB) is **Paget's disease of the breast**. It was first described by the British pathologist, James Paget in 1874. He described 14 cases of breast cancer associated with an eczema-like skin change of the nipple and areola. Almost 1% of all breast cancers present with Paget's disease of the nipple, and the diagnosis is confirmed by histologically demonstrating the Paget cells of the affected epidermis. The underlying breast cancer can be best demonstrated by combining all breast imaging methods. Of these, breast MRI is the most sensitive, showing the presence and true extent of the underlying DAB, often before calcifications can be detected on the mammogram.






2021  
BREAST SEMINAR SERIES of MEI

**László Tabár, MD, FACR (Hon)**  
*Course Director*


Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

*László Tabár, MD*  
*Tibor Tot, MD, Peter B. Dean, MD*





## Ductal Adenocarcinoma of the Breast (DAB), Part 5

Fluid producing DAB subtypes associated with calcifications




In 3D



Fluid producing micropapillary breast cancer of ductal origin (DAB)



In 3D



Neoductgenesis in micropapillary breast cancer of ductal origin (DAB)


This volume describes the subtypes of breast cancers that arise in the major ducts, produce a viscous, proteinaceous fluid. Little or no necrosis is present. The calcifications formed within the fluid have characteristic, but deceptively benign appearance, although the malignancy may extend throughout an entire lobe. This book will help identify these deceptive cases through correlating the mammographic/ultrasound/MRI presentation with large / thick section (3D) histology.

*László Tabár, MD*  
*Tibor Tot, MD, Peter B. Dean, MD*




Bloody and serous nipple discharge


## Ductal Adenocarcinoma of the Breast (DAB), Part 6



In 3D




Fluid producing micropapillary breast cancer of ductal origin (DAB)



Spontaneous unilateral serous or bloody nipple discharge can be an alarming clinical symptom for the patient and also, it may cause considerable differential diagnostic problem for the radiologist. This volume of our 3D book series correlates the imaging findings (mammography / breast ultrasound / breast MRI) with large thin- and large thick section (sub-gross, 3D) histology in cases when the underlying cause of the discharge is fluid-producing breast cancer originating from the major ducts (DAB).



László Tabár, MD  
Tibor Tot, MD, Peter B. Dean, MD






**In 3D**



Breast cancer originating from the major ducts

### Ductal Adenocarcinoma of the Breast (DAB), Part 7

Architectural distortion on the mammogram without calcifications or nipple discharge






Mammographic-MRI-subgross (3D) histologic correlation of this extensive micropapillary cancer originating from the major ducts presenting as architectural distortion.

Architectural distortion on the mammogram without calcifications or nipple discharge


**In 3D**







There are two main groups of diffuse breast cancers presenting on the mammogram as large regions of architectural distortion; these account for about 25% of all breast cancers and tend to have a poor outcome: 1) **Neoductgenesis**, i.e. "duct forming invasive carcinoma", the topic of this volume, often erroneously diagnosed as "DCIS", and 2) **Diffusely infiltrating breast cancer**, the topic of Vol. XI.

This volume demonstrates the DAB subgroup where the unnaturally high concentration of abnormal, tumor-filled ducts results in an asymmetric density with architectural distortion on the mammogram and often causes a palpable "thickening". Detecting architectural distortion on the mammogram and diagnosing the underlying disease correctly is a challenge for the radiologist. Breast cancers originating from the major ducts (DAB) are characterized by the formation of new, duct-like structures through the process of Neoductgenesis.


László Tabár, MD  
Tibor Tot, MD, Peter B. Dean, MD  
Olga Puchkova, MD




### Diffusely infiltrating breast cancer, Part 1



**In 3D**



Stereoscopic subgross (3D) image pair of a diffusely infiltrating breast cancer



**In 3D**

Extensive diffusely infiltrating breast cancer: the dominant feature is the extreme amount of connective tissue with concave contours.

This volume describes a breast cancer subtype that is a substantial challenge for the entire breast cancer team. The clinical, imaging and outcome observations indicate that diffusely infiltrating breast cancer represents a very unusual breast malignancy, regardless of whether it is E-cadherin negative or positive. All aspects of the diffusely infiltrating breast cancer suggest that it may have a site of origin different from all other breast cancers. We propose that it originates from the mesenchymal stem cells/progenitors through a complex process of epithelial-mesenchymal transformation and predominantly mesenchymal-epithelial transformation. Control of this unusual malignancy requires new approaches to earlier detection and entirely new therapeutic innovations.





2021  
BREAST SEMINAR SERIES of MEI

**László Tabár, MD, FACR (Hon)**  
*Course Director*

Detection and Diagnosis of Breast Diseases  
Using the Multimodality Approach. An interactive course.

The mission of the Tabar Foundation is to support research and education to fight against breast cancer. Dr. Tabár's own photographs are now available as high-quality prints. All proceeds from your tax-deductible purchase will support young physicians who are learning how to detect breast cancer when it is still curable. Visit: [tabarfoundation.org](http://tabarfoundation.org)

